



MASSACHUSETTS

Blue Cross Blue Shield of Massachusetts is an Independent Licensee of the Blue Cross and Blue Shield Association

Medical Policy

Endovascular Therapies for Extracranial Vertebral Artery Disease

Table of Contents

- [Policy: Commercial](#)
- [Policy: Medicare](#)
- [Authorization Information](#)
- [Coding Information](#)
- [Description](#)
- [Policy History](#)
- [Information Pertaining to All Policies](#)
- [References](#)

Policy Number: 730

BCBSA Reference Number: 7.01.148 (For Plan internal use only)

Related Policies

- Carotid, Vertebral and Intracranial Artery Stent Placement with or without Angioplasty [#219](#)
- Endovascular Procedures for Intracranial Arterial Disease, [#323](#)
- Endovascular Therapies for Extracranial Vertebral Artery Disease, [#730](#)

Policy

Commercial Members: Managed Care (HMO and POS), PPO, and Indemnity

Endovascular therapy, including percutaneous transluminal angioplasty with or without stenting, is considered **INVESTIGATIONAL** for the management of extracranial vertebral artery disease.

Prior Authorization Information

Inpatient

- For services described in this policy, precertification/preauthorization **IS REQUIRED** for all products if the procedure is performed **inpatient**.

Outpatient

- For services described in this policy, see below for products where prior authorization **might be required** if the procedure is performed **outpatient**.

	Outpatient
Commercial Managed Care (HMO and POS)	This is not a covered service.
Commercial PPO and Indemnity	This is not a covered service.

CPT Codes / HCPCS Codes / ICD Codes

Inclusion or exclusion of a code does not constitute or imply member coverage or provider reimbursement. Please refer to the member's contract benefits in effect at the time of service to determine coverage or non-coverage as it applies to an individual member.

Providers should report all services using the most up-to-date industry-standard procedure, revenue, and diagnosis codes, including modifiers where applicable.

The following codes are included below for informational purposes only; this is not an all-inclusive list.

The following CPT codes are considered investigational for **Commercial Members: Managed Care (HMO and POS), PPO, Indemnity, Medicare HMO Blue and Medicare PPO Blue:**

CPT Codes

CPT codes:	Code Description
0075T	Transcatheter placement of extracranial vertebral artery stent(s), including radiologic supervision and interpretation, open or percutaneous; initial vessel
0076T	Transcatheter placement of extracranial vertebral artery stent(s), including radiologic supervision and interpretation, open or percutaneous; each additional vessel (List separately in addition to code for primary procedure)

Description

Vertebrobasilar Circulation Ischemia

Ischemia of the vertebrobasilar or posterior circulation accounts for about 20% of all strokes. Posterior circulation strokes may arise from occlusion of the innominate and subclavian arteries, the extracranial vertebral arteries, or the intracranial vertebral, basilar, or posterior cerebral arteries. Compared with carotid artery disease, relatively little is known about the true prevalence of specific causes of posterior circulation strokes, particularly the prevalence of vertebral artery disease. In a report from a stroke registry, Gulli et al (2013) estimated that, in 9% of cases, posterior circulation strokes are due to stenosis of the proximal vertebral artery.¹ Patients who experience strokes or transient ischemic attacks of the vertebrobasilar circulation face a 25% to 35% risk of stroke within the subsequent 5 years. In particular, the presence of vertebral artery stenosis increases the 90-day risk of recurrent stroke by about 4-fold.

Relevant Clinical Anatomy and Pathophysiology

Large artery disease of the posterior circulation may be due to atherosclerosis (stenosis), embolism, dissection, or aneurysms. In about a third of cases, posterior circulation strokes are due to stenosis of the extracranial vertebral arteries or the intracranial vertebral, basilar, and posterior cerebral arteries. The proximal portion of the vertebral artery in the neck is the most common location of atherosclerotic stenosis in the posterior circulation. Dissection of the extracranial or intracranial vertebral arteries may also cause posterior circulation ischemia. By contrast, posterior cerebral artery ischemic events are more likely to be secondary to embolism from more proximal vessels.

The vertebral artery is divided into 4 segments, V1 through V4, of which segments V1, V2, and V3 are extracranial. V1 originates at the subclavian artery and extends to the C5 or C6 vertebrae; V2 crosses the bony canal of the transverse foramina from C2 to C5; V3 starts as the artery exits the transverse foramina at C2 and ends as the vessel crosses the dura mater and becomes an intracranial vessel. The most proximal segment (V1) is the most common location for atherosclerotic occlusive disease to occur, while arterial dissections are most likely to involve the extracranial vertebral artery just before the vessel crosses the dura mater. Compared with the carotid circulation, the vertebral artery system is more likely to be associated with anatomic variants, including a unilateral artery.

Atherosclerotic disease of the vertebral artery is associated with conventional risk factors for cerebrovascular disease. However, risk factors and the underlying pathophysiology of vertebral artery dissection and aneurysms differ. Extracranial vertebral artery aneurysms and dissections are most often secondary to trauma, particularly those with excessive rotation, distraction, or flexion/extension, or iatrogenic injury, such as during cervical spine surgeries. Spontaneous vertebral artery dissections are rare, and in many cases are associated with connective tissue disorders, including Ehlers-Danlos syndrome type IV, Marfan syndrome, autosomal dominant polycystic kidney disease, and osteogenesis imperfecta type I.²

Management of Extracranial Vertebral Artery Disease

The optimal management of occlusive extracranial vertebral artery disease is not well-defined. Medical treatment with antiplatelet or anticoagulant medications is a mainstay of therapy to reduce stroke risk. Medical therapy also typically involves risk reduction for classical cardiovascular risk factors. However, no randomized trials have compared specific antiplatelet or anticoagulant regimens.

Surgical revascularization may be used for vertebral artery atherosclerotic disease, but open surgical repair is considered technically challenging due to poor access to the vessel origin. Surgical repair may involve vertebral endarterectomy, bypass grafting, or transposition of the vertebral artery, usually to the common or internal carotid artery. Moderately-sized, single-center case series of surgical vertebral artery repair from 2012 and 2013 have reported overall survival rates of 91% and 77% at 3 and 6 years postoperatively, respectively, and arterial patency rates of 80% after 1 year of follow-up.^{3,4} Surgical revascularization may be used when symptomatic vertebral artery stenosis is not responsive to medical therapy, particularly when bilateral vertebral artery stenosis is present or when unilateral stenosis is present in the presence of an occluded or hypoplastic contralateral vertebral artery. Surgical revascularization may also be considered in patients with concomitant symptomatic carotid and vertebral disease who do not have relief from vertebrobasilar ischemia after carotid revascularization.

The management of extracranial vertebral artery aneurysms or dissections is controversial due to uncertainty about the risk of thromboembolic events associated with aneurysms and dissections. Antiplatelet therapy is typically used; surgical repair, which may include vertebral bypass, external carotid autograft, and vertebral artery transposition to the internal carotid artery, or endovascular treatment with stent placement or coil embolization, may also be used.

Given the technical difficulties related to surgically accessing the extracranial vertebral artery, endovascular therapies have been investigated for extracranial vertebral artery disease. Endovascular therapy may consist of percutaneous transluminal angioplasty, with or without stent implantation.

Summary

Vertebral artery diseases, including atherosclerotic stenosis, dissections, and aneurysms, can lead to ischemia of the posterior cerebral circulation. Conventional management of extracranial vertebral artery diseases may include medical therapy (eg, antiplatelet or anticoagulant medications), medications to reduce atherosclerotic disease risk (eg, statins), and/or surgical revascularization. Endovascular therapies have been investigated as an alternative to conventional management.

For individuals who have extracranial vertebral artery stenosis who receive percutaneous transluminal angioplasty (PTA) with or without stent implantation, the evidence includes randomized controlled trials (RCTs) and noncomparative studies. Relevant outcomes are overall survival, symptoms, morbid events, and treatment-related mortality and morbidity. Two RCTs, the Vertebral Artery Ischaemia Stenting Trial (VIST) and the Vertebral Artery Stenting Trial (VAST), found no advantage for endovascular intervention compared with best medical therapy alone. Evidence from noncomparative studies has shown that vertebral artery stenting can be performed with high rates of technical success and low periprocedural morbidity and mortality, and that vessel patency can be achieved in a high percentage of cases. However, long-term follow-up has demonstrated high rates of in-stent stenosis. The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

For individuals who have extracranial vertebral artery aneurysm(s), dissection(s), or arteriovenous fistula(e) who receive PTA with stent implantation, the evidence includes small case series and reports. Relevant outcomes are overall survival, symptoms, morbid events, and treatment-related mortality and morbidity. The available evidence has indicated that endovascular therapy for extracranial vertebral artery disorders other than stenosis is feasible and may be associated with favorable outcomes. However, given the lack of data comparing endovascular therapies to alternatives, the evidence is insufficient to permit conclusions about the efficacy of endovascular therapy for extracranial vertebral artery aneurysms, dissections, or arteriovenous fistulae. The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

Policy History

Date	Action
7/2023	Annual policy review. Description, summary, and references updated. Policy statements unchanged.
6/2022	Annual policy review. Description, summary, and references updated. Policy statements unchanged.
6/2021	Annual policy review. Description, summary, and references updated. Policy statements unchanged.
1/2021	Medicare information removed. See MP #132 Medicare Advantage Management for local coverage determination and national coverage determination reference.
7/2020	Annual policy review. Description, summary, and references updated. Policy statements unchanged.
6/2019	Annual policy review. Description, summary, and references updated. Policy statements unchanged.
6/2018	Annual policy review. New references added. Summary clarified.
7/2016	Annual policy review. New references added.
7/2015	New medical policy describing investigational indications. Effective 7/1/2015.

Information Pertaining to All Blue Cross Blue Shield Medical Policies

Click on any of the following terms to access the relevant information:

[Medical Policy Terms of Use](#)

[Managed Care Guidelines](#)

[Indemnity/PPO Guidelines](#)

[Clinical Exception Process](#)

[Medical Technology Assessment Guidelines](#)

References

1. Gulli G, Marquardt L, Rothwell PM, et al. Stroke risk after posterior circulation stroke/transient ischemic attack and its relationship to site of vertebrobasilar stenosis: pooled data analysis from prospective studies. *Stroke*. Mar 2013; 44(3): 598-604. PMID 23386676
2. Morasch MD, Phade SV, Naughton P, et al. Primary extracranial vertebral artery aneurysms. *Ann Vasc Surg*. May 2013; 27(4): 418-23. PMID 23540677
3. Coleman DM, Obi A, Criado E, et al. Contemporary outcomes after distal vertebral reconstruction. *J Vasc Surg*. Jul 2013; 58(1): 152-7. PMID 23478503
4. Ramirez CA, Febrer G, Gaudric J, et al. Open repair of vertebral artery: a 7-year single-center report. *Ann Vasc Surg*. Jan 2012; 26(1): 79-85. PMID 22176877
5. Markus HS, Larsson SC, Kuker W, et al. Stenting for symptomatic vertebral artery stenosis: The Vertebral Artery Ischaemia Stenting Trial. *Neurology*. Sep 19 2017; 89(12): 1229-1236. PMID 28835400
6. Compter A, van der Worp HB, Schonewille WJ, et al. Stenting versus medical treatment in patients with symptomatic vertebral artery stenosis: a randomised open-label phase 2 trial. *Lancet Neurol*. Jun 2015; 14(6): 606-14. PMID 25908089
7. Markus HS, Harshfield EL, Compter A, et al. Stenting for symptomatic vertebral artery stenosis: a preplanned pooled individual patient data analysis. *Lancet Neurol*. Jul 2019; 18(7): 666-673. PMID 31130429
8. Lattanzi S, Brigo F, Di Napoli M, et al. Endovascular treatment of symptomatic vertebral artery stenosis: A systematic review and meta-analysis. *J Neurol Sci*. Aug 15 2018; 391: 48-53. PMID 30103970
9. Xu R, Zhang X, Liu S, et al. Percutaneous transluminal angioplasty and stenting for vertebral artery stenosis. *Cochrane Database Syst Rev*. May 17 2022; 5(5): CD013692. PMID 35579383
10. Markus HS, Larsson SC, Dennis J, et al. Vertebral artery stenting to prevent recurrent stroke in symptomatic vertebral artery stenosis: the VIST RCT. *Health Technol Assess*. Aug 2019; 23(41): 1-30. PMID 31422789

11. Kikuchi T, Ishii A, Nakahara I, et al. Japanese Registry of Neuroendovascular Therapy: extracranial steno-occlusive diseases except for internal carotid artery stenosis. *Neurol Med Chir (Tokyo)*. 2014; 54(1): 40-5. PMID 24257542
12. Sun X, Ma N, Wang B, et al. The long term results of vertebral artery ostium stenting in a single center. *J Neurointerv Surg*. Dec 2015; 7(12): 888-91. PMID 25332411
13. Mohammadian R, Sharifipour E, Mansourizadeh R, et al. Angioplasty and stenting of symptomatic vertebral artery stenosis. Clinical and angiographic follow-up of 206 cases from Northwest Iran. *Neuroradiol J*. Aug 2013; 26(4): 454-63. PMID 24007733
14. Hatano T, Tsukahara T, Miyakoshi A, et al. Stent placement for atherosclerotic stenosis of the vertebral artery ostium: angiographic and clinical outcomes in 117 consecutive patients. *Neurosurgery*. Jan 2011; 68(1): 108-16; discussion 116. PMID 21099720
15. Pham MH, Rahme RJ, Arnaout O, et al. Endovascular stenting of extracranial carotid and vertebral artery dissections: a systematic review of the literature. *Neurosurgery*. Apr 2011; 68(4): 856-66; discussion 866. PMID 21242839
16. Badve MS, Henderson RD, O'Sullivan JD, et al. Vertebrobasilar dissections: case series comparing patients with and without dissecting aneurysms. *J Clin Neurosci*. Nov 2014; 21(11): 2028-30. PMID 24913932
17. Kondo R, Ishihara S, Uemiya N, et al. Endovascular Treatment for Acute Ischaemic Stroke Caused by Vertebral Artery Dissection: A Report of Three Cases and Literature Review. *NMC Case Rep J*. 2021; 8(1): 817-825. PMID 35079554
18. Horowitz MB, Miller G, Meyer Y, et al. Use of intravascular stents in the treatment of internal carotid and extracranial vertebral artery pseudoaneurysms. *AJNR Am J Neuroradiol*. Apr 1996; 17(4): 693-6. PMID 8730189
19. Felber S, Henkes H, Weber W, et al. Treatment of extracranial and intracranial aneurysms and arteriovenous fistulae using stent grafts. *Neurosurgery*. Sep 2004; 55(3): 631-8; discussion 638-9. PMID 15335430
20. Herrera DA, Vargas SA, Dublin AB. Endovascular treatment of traumatic injuries of the vertebral artery. *AJNR Am J Neuroradiol*. Sep 2008; 29(8): 1585-9. PMID 18499790
21. Ambekar S, Sharma M, Smith D, et al. Successful treatment of iatrogenic vertebral pseudoaneurysm using pipeline embolization device. *Case Rep Vasc Med*. 2014; 2014: 341748. PMID 25276469
22. Jang HJ, Oh SY, Shim YS, et al. Endovascular treatment of symptomatic high-flow vertebral arteriovenous fistula as a complication after c1 screw insertion. *J Korean Neurosurg Soc*. Oct 2014; 56(4): 348-52. PMID 25371787
23. Shang EK, Fairman RM, Foley PJ, et al. Endovascular treatment of a symptomatic extracranial vertebral artery aneurysm. *J Vasc Surg*. Nov 2013; 58(5): 1391-3. PMID 23561429
24. Takahashi S, Katayama K, Tatsugawa T, et al. A successful hybrid repair for vertebral arteriovenous fistula with extracranial vertebral artery aneurysm. *Ann Vasc Surg*. Jan 2015; 29(1): 126.e5-8. PMID 25304908
25. Kernan WN, Ovbiagele B, Black HR, et al. Guidelines for the prevention of stroke in patients with stroke and transient ischemic attack: a guideline for healthcare professionals from the American Heart Association/American Stroke Association. *Stroke*. Jul 2014; 45(7): 2160-236. PMID 24788967
26. Kleindorfer DO, Towfighi A, Chaturvedi S, et al. 2021 Guideline for the Prevention of Stroke in Patients With Stroke and Transient Ischemic Attack: A Guideline From the American Heart Association/American Stroke Association. *Stroke*. Jul 2021; 52(7): e364-e467. PMID 34024117
27. Brott TG, Halperin JL, Abbara S, et al. 2011 ASA/ACCF/AHA/AANN/AANS/ACR/ASNR/CNS/SAIP/SCAI/SIR/SNIS/SVM/SVS guideline on the management of patients with extracranial carotid and vertebral artery disease. A report of the American College of Cardiology Foundation/American Heart Association Task Force on Practice Guidelines, and the American Stroke Association, American Association of Neuroscience Nurses, American Association of Neurological Surgeons, American College of Radiology, American Society of Neuroradiology, Congress of Neurological Surgeons, Society of Atherosclerosis Imaging and Prevention, Society for Cardiovascular Angiography and Interventions, Society of Interventional Radiology, Society of NeuroInterventional Surgery, Society for Vascular Medicine, and Society for Vascular Surgery. *Circulation*. Jul 26 2011; 124(4): e54-130. PMID 21282504

28. Centers for Medicare & Medicaid Services. National Coverage Determination (NCD) for Percutaneous Transluminal Angioplasty (PTA) (20.7). 2013; <https://www.cms.gov/medicare-coverage-database/details/ncd-details.aspx?NCDId=201>. Accessed March 16, 2023.